

D. S. SILVESTROV AND M. O. DROZDENKO

NECESSARY AND SUFFICIENT CONDITIONS FOR WEAK CONVERGENCE OF FIRST-RARE-EVENT TIMES FOR SEMI-MARKOV PROCESSES. II

Necessary and sufficient conditions for weak convergence of first-rare-event times for semi-Markov processes, obtained in the first part of this paper [66], are applied to counting processes generating by flows of rare events controlled by semi-Markov processes, random geometric sums, and risk processes. In particular, necessary and sufficient conditions for stable approximation of ruin probabilities including the case of diffusion approximation are given.

4. INTRODUCTION

In the first part of the present paper [66], necessary and sufficient conditions for weak convergence of first-rare-event times for semi-Markov processes. In the second part of the paper, we apply these results to counting processes generating by flows of rare events controlled by semi-Markov processes, random geometric sums, and risk processes. In particular, we give necessary and sufficient conditions for stable approximation of ruin probabilities including the case of diffusion approximation are given.

We use all notations and conditions introduced in the first part of the present paper as well as continue numbering of sections, theorems and lemmas began there.

First-rare-event times for semi-Markov processes considered in the first part of the present paper reduce to random geometric sums in the case of degenerated imbedded Markov chain.

In this way, our results are connected with asymptotical results for geometric sums. Here, we would like first to mention originating papers by Rényi (1956), who first formulated conditions of convergence of geometrical sums to exponential law, and works by Kovalenko (1965), Gnedenko and Fraier (1969), Szynal (1976), Korolev (1989), Kalashnikov and Vsekhsvyatski (1989), Melamed (1989), Kruglov and Korolev (1990), Kartashov (1991), Gnedenko and Korolev (1996), Kalashnikov (1997), Bening and Korolev (2002).

It should be noted that the class of possible limiting laws for standard geometric random sums, with summands independent on geometric random

2000 *Mathematics Subject Classification*. Primary 60K15, 60F17, 60K20.

Key words and phrases. Weak convergence, semi-Markov processes, first-rare-event time, limit theorems, necessary and sufficient conditions.

indices and possessing regularly varying tail probabilities, was described by Kovalenko (1965) who also credited for necessary and sufficient conditions for weak convergence of such sums. These results were generalized by Kruglov and Korolev (1990), in particular, to the case of triangular array mode. Our results yield necessary and sufficient conditions for weak convergence summands have regularly varying tail probabilities but for more general geometric sums with summands that can depend on geometric random indices via indicators of rare events. Also, our conditions have different and, as we think, more convenient form than in the works mentioned above.

We give also necessary and sufficient conditions for weak convergence of counting processes generating by the corresponding flows of rare events to standard renewal flows. These results are connected with the results related to studies of convergence for rareficated stochastic flows given in Rényi (1956), Belyaev (1963), Kovalenko (1965), Mogyoródi (1971, 1972a, 1972b), Szantai (1971a, 1971b), Råde (1972a, 1972b), Zakusilo (1972a, 1972b), Jagers (1974), Jagers and Lindvall (1974), Tomko (1974), Kallenberg (1975) and Lindvall (1976), Serfozo (1976, 1984a, 1984b), Gasanenko (1980), and Böker and Serfozo (1983).

As possible areas of applications, we can point out limit theorems for different lifetime functionals such as occupation times or waiting times in queuing systems, lifetime in reliability models, extinction times in population dynamic models, ruin times for insurance models, etc. We mention here works by Gnedenko (1964a, 1964b), Gnedenko and Kovalenko (1964, 1987), Silvestrov (1969), Masol (1973), Kovalenko (1975, 1977, 1980, 1994), Solov'ev (1971, 1983), Ivchenko, Kashtanov, and Kovalenko (1979), Kovalenko and Kuznetsov (1988), Kovalenko, Kuznetsov, Pegg (1997), Englund (1999a, 1999b), Anisimov, Zakusilo, and Donchenko (1987), Assmussen (1987, 2003), Kalashnikov and Rachev (1990), Gyllenberg and Silvestrov (1994).

We apply our results to asymptotical analysis of non-ruin probabilities for risk processes. We study asymptotics for non-ruin probabilities in the critical case when the initial capital of an insurance company tends to infinity and simultaneously the safety loading coefficient tends to 1. Two cases are considered, when the claim distribution belongs to the domain of attraction of a degenerated or a stable law. The first case corresponds to the classical model of diffusion approximation, the second one can be referred as the case of stable approximation for non-ruin probabilities.

As was mention above, we give necessary and sufficient conditions for stable approximation for non-ruin probabilities, including the case of diffusion approximation.

Our results are connected with the results related to various sufficient conditions for diffusion approximation given in Iglehart (1969), Siegmund (1975), Harrison (1977), Grandell (1977, 1991), Gerber (1979), Asmussen

(1987, 1989, 2000, 2003), Glynn (1990), Schmidli (1992, 1997), Gyllenberg and Silvestrov (1999, 2000b), and Silvestrov (2000b).

It is not out of picture to note that asymptotical relations related to diffusion approximation for ruin probabilities can be also interpreted in terms of the queue theory as variants of so-called heavy traffic approximation for waiting times (see, for instance, Asmussen (1987)).

The results presented in the paper are also connected with results related to asymptotics of ruin probabilities for risk processes with heavy tailed claim distributions given by von Bahr (1975), Thorin and Wikstad (1976), Embrechts, Goldie and Veraverbeke (1979), Embrechts and Veraverbeke (1982), Asmussen (1996, 2000, 2003), Embrechts (1997), Klüppelberg and Stadmular (1998), Zinchenko (1999). It should be noted, however, that the latter group of works relates to so-called sub-critical case when the safety loading coefficient less than and separated of 1.

The paper is organized in the following way. In Section 2 we give necessary and sufficient of weak convergence of counting processes generating by the corresponding flows of rare events. As was mentioned above, the model of first-rare-event times for semi-Markov processes reduces to the model of geometrical random sums in the case of degenerated imbedded Markov chain. In Section 3, we present applications of results obtained in the first part of the present paper to random geometric sums as well as give necessary and sufficient conditions for stable approximation for non-ruin probabilities.

5. FLOWS OF RARE EVENTS

In this section we study conditions of convergence for the flows of rare events connected with the model considered above.

Let us define recursively random variables

$$\nu_\varepsilon(k) = \min(n \geq \nu_\varepsilon(k-1) : \zeta_n \in D_\varepsilon), \quad k = 1, 2, \dots,$$

where $\nu_\varepsilon(0) = 0$. A random variable $\nu_\varepsilon(k)$ counts the number of transitions of the imbedded Markov chain η_n up to the k -th appearance of the “rare” event $\{\zeta_n \in D_\varepsilon\}$. Obviously, $\nu_\varepsilon(1) = \nu_\varepsilon$.

Let us also define inter-rare-event times,

$$\varkappa_\varepsilon(k) = \sum_{n=\nu_\varepsilon(k-1)}^{\nu_\varepsilon(k)} \varkappa_n, \quad k = 1, 2, \dots$$

Let us also introduce random variables showing positions of the imbedded Markov chain η_n at moments $\nu_\varepsilon(k)$,

$$\eta_\varepsilon(k) = \eta_{\nu_\varepsilon(k)}, \quad k = 0, 1, \dots$$

Obviously $(\eta_\varepsilon(k), \varkappa_\varepsilon(k)), k = 0, 1, \dots$ (here $\varkappa_\varepsilon(k) = 0$) is a Markov renewal process, i.e. a homogeneous Markov chain with the phase space

$X \times [0, \infty)$ and transition probabilities,

$$\begin{aligned}
 (1) \quad & \mathbb{P}\{\eta_\varepsilon(k+1) = j, \varkappa_\varepsilon(k+1) \leq t/\eta_\varepsilon(k) = i, \varkappa_\varepsilon(k) = s\} \\
 & = \mathbb{P}\{\eta_\varepsilon(k+1) = j, \varkappa_\varepsilon(k+1) \leq t/\eta_\varepsilon(k) = i\} \\
 & = Q_{ij}^{(\varepsilon)}(t) = \mathbb{P}_i\{\eta_{\nu_\varepsilon} = j, \xi_\varepsilon \leq t\}, \quad i, j \in X, \quad s, t \geq 0.
 \end{aligned}$$

Let us now define random variables,

$$\xi_\varepsilon(k) = \sum_{n=1}^{\nu_\varepsilon(k)} \varkappa_n = \sum_{n=1}^k \varkappa_\varepsilon(n), \quad k = 0, 1, \dots$$

Random variable $\xi_\varepsilon(k)$ can be interpreted as the time of k -th appearance of the rare event time for the semi-Markov process $\eta(t)$. Obviously, $\xi_\varepsilon(0) = 0$ and $\xi_\varepsilon(1) = \xi_\varepsilon$.

Now we can define a counting stochastic process that describes the flow of rare events,

$$N_\varepsilon(t) = \max(k \geq 0 : \xi_\varepsilon(k) \leq tu_\varepsilon), \quad t \geq 0.$$

Note that the time scale for this counting process is stretched with the use of the scale parameter u_ε according the asymptotic results given in Theorem 1.

Let us also define the corresponding limiting counting process. Let $\varkappa(k)$, $k = 1, 2, \dots$ be a sequence of positive i.i.d. random variables with the distribution $F_{a,\gamma}(u)$, and

$$\xi(k) = \sum_{n=1}^k \varkappa(n), \quad n = 0, 1, \dots$$

Let us also define the standard renewal counting process with i.i.d. inter-renewal times $\varkappa(k)$, $k = 1, 2, \dots$,

$$N(t) = \max(k \geq 0 : \xi(k) \leq t), \quad t \geq 0.$$

Theorem 2. *Let conditions **A**, **B**, **C**, and **D** hold. Then, the class of all possible non-zero limiting counting processes (in the sense of weak convergence of finite-dimensional distributions) for the counting processes $N_\varepsilon(t), t \geq 0$ coincides with the class of standard renewal counting process $N(t), t \geq 0$ with the distribution function of inter-renewal time $F_{a,\gamma}(u)$ where $0 < \gamma \leq 1, a > 0$. Conditions \mathbf{E}_γ and $\mathbf{F}_{a,\gamma}$ are necessary and sufficient for such convergence in the case when the corresponding limiting counting process has the distribution function of inter-renewal time $F_{a,\gamma}(u)$.*

Proof. Obviously,

$$F_i^{(\varepsilon)}(u) = \mathbb{P}_i\{\xi_\varepsilon \leq u\} = \sum_{j \in X} Q_{ij}^{(\varepsilon)}(t), \quad u \geq 0.$$

Using Markov property of the Markov renewal process $(\eta_\varepsilon(k), \varkappa_\varepsilon(k))$ we get the following formula for joint distributions of the properly normalized inter-renewal times for the counting process $N_\varepsilon(t)$,

$$(2) \quad \begin{aligned} & \mathbf{P}_i\{\varkappa_\varepsilon(k)/u_\varepsilon \leq t_k, k = 1, \dots, n\} \\ &= \sum_{j \in X} \mathbf{P}_i\{\varkappa_\varepsilon(k)/u_\varepsilon \leq t_k, k = 1, \dots, n-1, \eta_\varepsilon(n-1) = j\} \\ & \times F_j^{(\varepsilon)}(t_n u_\varepsilon), \quad i \in X, t_1, \dots, t_n \geq 0, \quad n = 1, 2, \dots \end{aligned}$$

According Theorem 1, under **A**, **B**, **C**, and **D**, conditions \mathbf{E}_γ and $\mathbf{F}_{a,\gamma}$ imply that (a) $F_j^{(\varepsilon)}(t_n u_\varepsilon) \rightarrow F_{a,\gamma}(t_n)$ as $\varepsilon \rightarrow 0, t_n \geq 0, j \in X$.

Using (a) and relation (2) we get that, under **A**, **B**, **C**, and **D**, conditions \mathbf{E}_γ and $\mathbf{F}_{a,\gamma}$ imply that, for every $i \in X, n = 1, 2, \dots, t_1, \dots, t_n \geq 0$,

$$(3) \quad \begin{aligned} & \mathbf{P}_i\{\varkappa_\varepsilon(k)/u_\varepsilon \leq t_k, k = 1, \dots, n\} \\ & \rightarrow \prod_{k=1}^n F_{a,\gamma}(t_k) \text{ as } \varepsilon \rightarrow 0. \end{aligned}$$

Relation (3) means that the inter-renewal times $\varkappa_\varepsilon(k)/u_\varepsilon, k = 1, 2, \dots$ are asymptotically independent. Note that the multivariate distribution function on the right hand side in (3) is continuous. Due to this fact, relation (3) implies in an obvious way that, for every $i \in X$, real $0 = t_0 \leq t_1 \leq \dots \leq t_n$, integer $0 \leq r_1 \leq \dots \leq r_n, n = 1, 2, \dots$,

$$(4) \quad \begin{aligned} & \mathbf{P}_i\{N_\varepsilon(t_k) \geq r_k, k = 1, \dots, n\} \\ &= \mathbf{P}_i\{\xi_\varepsilon(r_k)/u_\varepsilon \leq t_k, k = 1, \dots, n\} \\ & \rightarrow \mathbf{P}_i\{\xi(r_k) \leq t_k, k = 1, \dots, n\} \\ &= \mathbf{P}_i\{N(t_k) \geq r_k, k = 1, \dots, n\} \text{ as } \varepsilon \rightarrow 0, \end{aligned}$$

The statement of necessity follows is trivial and follows from the following formula

$$\mathbf{P}_i\{N_\varepsilon(t) \geq 1\} = \mathbf{P}_i\{\xi_\varepsilon/u_\varepsilon \leq t\}, \quad t \geq 0.$$

The proof is complete. \square

What is interesting, that under **A**, **B**, **C**, and **D**, conditions \mathbf{E}_γ and $\mathbf{F}_{a,\gamma}$ are not sufficient for weak convergence of transition probabilities of the Markov renewal process $(\eta_\varepsilon(k), \varkappa_\varepsilon(k))$ that forms the counting process $N_\varepsilon(t)$.

It follows from the following lemma which describes the asymptotic behavior of so-called ‘‘absorbing’’ probabilities,

$$Q_{ij}^{(\varepsilon)}(\infty) = \mathbf{P}_i\{\eta_{\nu_\varepsilon} = j\}, \quad i, j \in X.$$

Let us denote

$$p_{i\varepsilon}(r) = \mathbf{P}_i\{\zeta_1 \in D_\varepsilon, \eta_1 = r\}, \quad i, r \in X,$$

and

$$p_\varepsilon(r) = \sum_{i=1}^m \pi_i p_{i\varepsilon}(r), \quad j \in X.$$

By the definition,

$$(5) \quad p_\varepsilon = \sum_{i \in X} \pi_i \mathbf{P}_i \{ \zeta_1 \in D_\varepsilon \} = \sum_{r \in X} p_\varepsilon(r).$$

Lemma 7. *Let conditions **B**, **C** hold. Then, for every $i \in X$,*

$$(6) \quad Q_{ir}^{(\varepsilon)}(\infty) - \frac{p_\varepsilon(r)}{p_\varepsilon} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0, \quad r \in X.$$

Proof. Let us define the probability that the first rare event will occur when the state of the imbedded Markov chain will be r and before the first hitting of the imbedded Markov chain in the state i , under condition that the initial state of this Markov chain $\eta_0 = j$,

$$q_{ji\varepsilon}(r) = \mathbf{P}_j \{ \nu_\varepsilon \leq \tau_i, \eta_{\nu_\varepsilon} = r \}, \quad i, j, r \in X.$$

Taking into account that the Markov renewal process $(\eta_n, \mathcal{Z}_n, \zeta_n)$ regenerates at moments of return to every state i and ν_ε is a Markov moment for this process, we can get following cyclic representation for absorbing probabilities $Q_{ir}^{(\varepsilon)}(\infty)$,

$$(7) \quad \begin{aligned} Q_{ir}^{(\varepsilon)}(\infty) &= \sum_{n=0}^{\infty} \mathbf{P}_i \{ \tau_i(n) < \nu_\varepsilon \leq \tau_i(n+1), \eta_{\nu_\varepsilon} = j \} \\ &= \sum_{n=0}^{\infty} (1 - q_{i\varepsilon})^n q_{ii\varepsilon}(r) = \frac{q_{ii\varepsilon}(r)}{q_{i\varepsilon}}, \quad i, r \in X. \end{aligned}$$

The probabilities $q_{ji\varepsilon}(r), j \in X$ satisfy, for every $i, r \in X$, the following system of linear equations similar with system (20) given in the proof of Lemma 1,

$$(8) \quad \begin{cases} q_{ji\varepsilon}(r) = p_{j\varepsilon}(r) + \sum_{k \neq i} p_{jk}^{(\varepsilon)} q_{ki\varepsilon}(r) \\ j \in X \end{cases}$$

This system has the matrix of coefficients ${}_i \mathbf{P}^{(\varepsilon)}$ as the system of linear equations (20) mentioned above and differs of this system only by the free terms. Thus by repeating reasoning given in the proof of Lemma 1 we can get the following formula similar with formula (24) also given in the proof of Lemma 1,

$$(9) \quad q_{ii\varepsilon}(r) = \sum_{k=1}^m \mathbf{E}_i \delta_{ik\varepsilon} p_{k\varepsilon}(r).$$

Recall that it was shown in the proof of Lemma 1 that **(b)** $\mathbf{E}_i \delta_{ik\varepsilon} \rightarrow \pi_k / \pi_i$ as $\varepsilon \rightarrow 0$, for $i, k \in X$. Using **(b)**, relation **(c)** $\pi_i q_{i\varepsilon} / p_\varepsilon \rightarrow 1$ given in

Lemma 1, and inequality **(d)** $p_\varepsilon(r) \leq p_\varepsilon$, following from formula (5), we get, for every $i, r \in X$,

$$(10) \quad \begin{aligned} \frac{\left| q_{ii\varepsilon}(r) - \frac{p_\varepsilon(r)}{\pi_i} \right|}{q_{i\varepsilon}} &\leq \sum_{k=1}^m \left| \mathbb{E}_i \delta_{ik\varepsilon} - \frac{\pi_k}{\pi_i} \right| \cdot \frac{\pi_i p_{k\varepsilon}(r)}{\sum_{j=1}^m \pi_j p_{j\varepsilon}} \cdot \frac{p_\varepsilon}{\pi_i q_{i\varepsilon}} \\ &\leq \sum_{k=1}^m \left| \mathbb{E}_i \delta_{ik\varepsilon} - \frac{\pi_k}{\pi_i} \right| \cdot \frac{\pi_i}{\pi_k} \cdot \frac{p_\varepsilon}{\pi_i q_{i\varepsilon}} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \end{aligned}$$

Using **(c)** and **(d)** ones more time we get,

$$(11) \quad \begin{aligned} \left| \frac{p_\varepsilon(r)}{\pi_i q_{i\varepsilon}} - \frac{p_\varepsilon(r)}{p_\varepsilon} \right| &\leq \frac{p_\varepsilon(r)}{p_\varepsilon} \cdot \frac{\left| q_{i\varepsilon} - \frac{p_\varepsilon}{\pi_i} \right|}{\pi_i q_{i\varepsilon}} \\ &\leq \frac{\left| q_{i\varepsilon} - \frac{p_\varepsilon}{\pi_i} \right|}{p_\varepsilon} \cdot \frac{p_\varepsilon}{\pi_i q_{i\varepsilon}} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \end{aligned}$$

Formula (7) together with relations (10) and (11) imply in an obvious way relation (6). This completes the proof. \square

Let us introduce the following balancing condition:

$$\mathbf{L}:: \frac{p_\varepsilon(j)}{p_\varepsilon} \rightarrow Q_j \text{ as } \varepsilon \rightarrow 0, \quad j \in X.$$

Constants Q_j , automatically satisfy the following conditions **(e₁)** $Q_j \geq 0, j \in X$, and **(e₂)** $\sum_{j \in X} Q_j = 1$.

Lemma 7 implies the following statement.

Lemma 8. *Let conditions **B**, **C** hold. Then, condition **L** is necessary and sufficient for the following relation to hold (for some or every $i \in X$, respectively, in the statements of necessity and sufficiency),*

$$(12) \quad Q_{ir}^{(\varepsilon)}(\infty) \rightarrow Q_r \text{ as } \varepsilon \rightarrow 0, \quad r \in X.$$

The following theorem shows that the-first-rare-event times ξ_ε and random functional η_{ν_ε} are asymptotically independent, and completes the description of the asymptotic behavior of the transition probabilities $Q_{ij}^{(\varepsilon)}(t)$ for the Markov renewal process $(\eta_\varepsilon(k), \varkappa_\varepsilon(k))$.

Theorem 3. *Let conditions **A**, **B**, **C**, and **D** hold. Then, conditions **E_γ**, **F_{a,γ}** and **L** are necessary and sufficient for the asymptotic relation (2) given in Theorem 1 and the asymptotic relation (12) given in Lemma 8 to hold. In this case, for every $u \in [0, \infty], i, r \in X$,*

$$(13) \quad Q_{ir}^{(\varepsilon)}(uu_\varepsilon) \rightarrow F_{a,\gamma}(u)Q_r \text{ as } \varepsilon \rightarrow 0.$$

Proof. The first statement of the theorem follows from Theorem 1 and Lemma 8. Let us prove that conditions \mathbf{E}_γ , $\mathbf{F}_{a,\gamma}$ and \mathbf{L} imply the asymptotic relation (13).

Let us introduce, for $i, r \in X$, Laplace transforms,

$$\Phi_{ir\varepsilon}(s) = \mathbf{E}_i \exp\{-s\xi_\varepsilon\} \chi(\eta_{\nu_\varepsilon} = r), \quad s \geq 0,$$

and

$$\tilde{\psi}_{ir\varepsilon}(s) = \mathbf{E}_i \{\exp\{-s\tilde{\beta}_{i\varepsilon}\} \chi(\eta_{\nu_\varepsilon} = r) / \nu_\varepsilon \leq \tau_i\}, \quad s \geq 0.$$

Analogously to formula (4) given in the proof of Theorem 1 the following representation can be written down for Laplace transforms $\Phi_{ir\varepsilon}(s)$,

$$\begin{aligned} \Phi_{ir\varepsilon}(s) &= \sum_{n=0}^{\infty} (1 - q_{i\varepsilon})^n q_{i\varepsilon} \psi_{i\varepsilon}(s)^n \tilde{\psi}_{ir\varepsilon}(s) \\ &= \frac{q_{i\varepsilon} \tilde{\psi}_{ir\varepsilon}(s)}{1 - (1 - q_{i\varepsilon}) \psi_{i\varepsilon}(s)} \\ (14) \quad &= \frac{1}{1 + (1 - q_{i\varepsilon}) \frac{(1 - \psi_{i\varepsilon}(s))}{q_{i\varepsilon}}} \cdot \tilde{\psi}_{ir\varepsilon}(s), \quad s \geq 0. \end{aligned}$$

Let us now show that under, conditions \mathbf{A} , \mathbf{B} , and \mathbf{C} , for every $s \geq 0$ and $i, r \in X$,

$$(15) \quad \tilde{\psi}_{ir\varepsilon}(s/u_\varepsilon) - q_{ii\varepsilon}(r)/q_{i\varepsilon} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Indeed, using Lemma 2 we get, for any $\delta > 0$,

$$\begin{aligned} &\frac{q_{ii\varepsilon}(r)}{q_{i\varepsilon}} - \tilde{\psi}_{ir\varepsilon}\left(\frac{s}{u_\varepsilon}\right) \\ (16) \quad &= \mathbf{E}_i \left\{ (1 - \exp\{-s\frac{\tilde{\beta}_{i\varepsilon}}{u_\varepsilon}\}) \chi(\eta_{\nu_\varepsilon} = r) / \nu_\varepsilon \leq \tau_i \right\} \\ &\leq (1 - e^{s\delta}) + \mathbf{P}_i \left\{ \frac{\tilde{\beta}_{i\varepsilon}}{u_\varepsilon} \geq \delta / \nu_\varepsilon \leq \tau_i \right\} \rightarrow (1 - e^{s\delta}) \text{ as } \varepsilon \rightarrow 0. \end{aligned}$$

Relation (16) implies, due to possibility of an arbitrary choice of $\delta > 0$, relation (15).

Using formula (14) and relation (29), given in Lemma 2, and (15), Theorem 1 and Lemma 8 we get, for every $s \geq 0$ and $i, r \in X$,

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \Phi_{ir\varepsilon}(s/u_\varepsilon) \\ (17) \quad &= \lim_{\varepsilon \rightarrow 0} \frac{1}{1 + (1 - q_{i\varepsilon}) \frac{(1 - \psi_{i\varepsilon}(s))}{q_{i\varepsilon}}} \cdot \tilde{\psi}_{ir\varepsilon}(s/u_\varepsilon) \\ &= \lim_{\varepsilon \rightarrow 0} \Phi_{i\varepsilon}(s/u_\varepsilon) \cdot Q_{ir}^{(\varepsilon)}(\infty) \\ &= \frac{1}{1 + as^\gamma} \cdot Q_r. \end{aligned}$$

Relation (17) equivalent to relation (13). \square

6. GEOMETRIC SUMS AND STABLE APPROXIMATION
FOR NON-RUIN PROBABILITIES

In this section we apply our results to so-called geometric random sums. This is reduction of our model for the case when the imbedded Markov chain η_n has the degenerated set of states $X = \{1\}$.

In this case, the first-rare-event time $\xi_\varepsilon = \sum_{n=1}^{\nu_\varepsilon} \varkappa_n$ is a geometric sum.

Indeed, $(\varkappa_n, \zeta_n), n = 1, 2, \dots$ is a sequence of i.i.d. random vectors. Therefore the random variable

$$\nu_\varepsilon = \min(n \geq 1 : \zeta_n \in D_\varepsilon)$$

has a geometric distribution with the success probability

$$p_\varepsilon = \mathbf{P}\{\zeta_n \in D_\varepsilon\}.$$

However, the geometric random index ν_ε and random summands $\varkappa_n, n = 1, 2, \dots$ are, in this case, dependent random variables. They depend via the indicators of rare events $\chi_{n\varepsilon} = \chi(\zeta_n \in D_\varepsilon), n = 1, 2, \dots$. More precisely, $(\varkappa_n, \chi_{n\varepsilon}, n = 1, 2, \dots)$ is a sequence of i.i.d. random vectors.

Conditions **A** and **B** take, in this case, the following form:

$$\mathbf{A}':: \lim_{t \rightarrow \infty} \overline{\lim}_{\varepsilon \rightarrow 0} \mathbf{P}\{\varkappa_1 > t/\zeta_1 \in D_\varepsilon\} = 0;$$

and

$$\mathbf{B}':: 0 < p_\varepsilon = \mathbf{P}\{\zeta_1 \in D_\varepsilon\} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Condition **C** holds automatically.

Condition **D** remains and should be imposed on the function p_ε^{-1} , defined in condition **B'**, and the normalization function u_ε .

Conditions **E** $_\gamma$ and **F** $_{a,\gamma}$ remain and should be imposed on the distribution function $G(t) = \mathbf{P}\{\varkappa_1 \leq t\}$ (no averaging is involved).

A standard geometric sum is particular case of the model described above, which corresponds to the case, when two sequences of random variables $\varkappa_n, n = 1, 2, \dots$ and $\zeta_n, n = 1, 2, \dots$ are independent. In this case, the random index ν_ε and summands $\varkappa_n, n = 1, 2, \dots$ are also independent.

Note that a standard geometric sum with any distribution of summands $G(t)$ and parameter of geometric random index $p_\varepsilon \in (0, 1]$ can be modelled in this way. Indeed, it is enough to consider the geometric sum

$$\xi_\varepsilon = \sum_{n=1}^{\nu_\varepsilon} \varkappa_n$$

defined above, where **(a₁)** $\varkappa_n, n = 1, 2, \dots$ is a sequence of i.i.d. random variables with the distribution function $G(t)$;

$$\mathbf{(a_2)} \quad \nu_\varepsilon = \max(n \geq 1 : \zeta_n \in D_\varepsilon),$$

where $\zeta_n, n = 1, 2, \dots$ is a sequence of i.i.d. random variables uniformly distributed in the interval $[0, 1]$ and domains $D_\varepsilon = [0, p_\varepsilon]$; **(a₃)** two sequences of random variables $\varkappa_n, n = 1, 2, \dots$ and $\zeta_n, n = 1, 2, \dots$ are independent.

In the case of standard geometric sums, condition **A** holds automatically. Theorem 1 reduces in this case to the result equivalent to those obtained by Kovalenko (1965). The difference is in the form of necessary and sufficient conditions. Conditions \mathbf{G}_γ and $\mathbf{H}_{a,\gamma}$, based on Laplace transforms of distributions $G(t)$, were used in this work. Our results are based on conditions \mathbf{E}_γ and $\mathbf{F}_{a,\gamma}$, based on distributions $G(t)$, and have, as we think, a more transparent form.

Let us illustrate applications of Theorem 1 by giving necessary and sufficient conditions for stable approximation of non-ruin probabilities. Let us consider a process used in classical risk theory to model the business of an insurance company,

$$X_\varepsilon(t) = c_\varepsilon t - \sum_{n=1}^{N_\lambda(t)} Z_n, \quad t \geq 0.$$

Here, a positive constant c_ε (depending on parameter $\varepsilon > 0$) is the gross premium rate, $N_\lambda(t)$, $t \geq 0$ is a Poissonian process with parameter λ counting the number of claims on an insurance company in the time-interval $[0, t]$, and Z_n , $n = 1, 2, \dots$ is a sequence of nonnegative i.i.d. random variables, which are independent on the process $N_\lambda(t)$, $t \geq 0$. The random variable Z_k is the amount of the k^{th} claim.

An important object for studies in this model is the non-ruin probabilities on infinite time interval for a company with an initial capital $u \geq 0$,

$$F_\varepsilon(u) = \mathbf{P} \left\{ u + \inf_{t \geq 0} X(t) \geq 0 \right\}, \quad u \geq 0.$$

Let $H(x) = \mathbf{P}\{Z_1 \leq x\}$ be a claim distribution function. We assume the standard condition:

$$\mathbf{M}:: \mu = \int_0^\infty sH(ds) < \infty.$$

The crucial role in is plaid by the so-called safety loading coefficient

$$\alpha_\varepsilon = \lambda\mu/c_\varepsilon.$$

If $\alpha_\varepsilon \geq 1$ then $F_\varepsilon(u) = 0$, $u \geq 0$. The only non-trivial case is when $\alpha_\varepsilon < 1$. We assume the following condition:

$$\mathbf{N}:: \alpha_\varepsilon < 1 \text{ for } \varepsilon > 0 \text{ and } \alpha_\varepsilon \rightarrow 1 \text{ as } \varepsilon \rightarrow 0.$$

According to Pollaczek–Khinchine formula (see, for example, Asmussen (2000)), the non-ruin distribution function $F_\varepsilon(u)$ coincides with distribution function of a geometric random sum which is slightly differ on the standard geometric sums considered above. Namely,

$$(18) \quad F_\varepsilon(u) = \mathbf{P}\{\xi'_\varepsilon = \sum_{n=1}^{\nu_\varepsilon-1} \varkappa_n \leq u\}, \quad u \geq 0,$$

where (a) $\varkappa_n, n = 1, 2, \dots$ is a sequence of non-negative i.i.d. random variables with distribution function

$$G(u) = \frac{1}{\mu} \int_0^u (1 - H(s)) ds, \quad u \geq 0$$

(so-called steady claim distribution);

(b) $\nu_\varepsilon = \min(n \geq 1, \quad \chi_{n\varepsilon} = 1);$

and (c) $\chi_{n\varepsilon}, n = 1, 2, \dots,$ is a sequence of i.i.d. random variables taking values 1 and 0 with probabilities $p_\varepsilon = 1 - \alpha_\varepsilon$ and $1 - p_\varepsilon$; (d) random sequences $\varkappa_n, n = 1, 2, \dots$ and $\chi_{n\varepsilon}, n = 1, 2, \dots$ are independent.

As was mentioned above condition **A** holds automatically in this case, and, therefore, due to Remark 12, Theorem 1, which specification to the geometric sums was described above, can be applied to the geometric random sums ξ'_ε .

Conditions **A** and **C** can be omitted. Condition **B** is equivalent to condition **N**. Condition **D** takes the following form:

$$\mathbf{D}':: u_\varepsilon, p_\varepsilon^{-1} = (1 - \alpha_\varepsilon)^{-1} \in \mathbb{W}.$$

Conditions \mathbf{E}_γ and $\mathbf{F}_{a,\gamma}$ ($a > 0$ and $0 < \gamma \leq 1$) take, in this case, the following form:

$$\mathbf{E}'_\gamma:: \frac{t \int_t^\infty (1-H(s)) ds}{\int_0^t s(1-H(s)) ds} \rightarrow \frac{1-\gamma}{\gamma} \text{ as } t \rightarrow \infty.$$

$$\mathbf{F}'_{a,\gamma}:: \frac{\int_0^{u_\varepsilon} s(1-H(s)) ds}{(1-\alpha_\varepsilon)\mu u_\varepsilon} \rightarrow a \frac{\gamma}{\Gamma(2-\gamma)} \text{ as } \varepsilon \rightarrow 0.$$

Let us summarize the discussion above in the form of the following theorem which gives necessary and sufficient conditions for stable approximation of non-ruin probabilities.

Theorem 4. *Let conditions **M**, **N**, and **D'** hold. Then the class of all possible non-concentrated in zero limiting (in the sense of weak convergence) distribution functions $F(u)$, such that the non-ruin distribution functions $F_\varepsilon(uu_\varepsilon) \Rightarrow F(u)$ as $\varepsilon \rightarrow 0$, coincides with the class of distributions $F_{a,\gamma}(u)$ with Laplace transforms $\frac{1}{1+as^\gamma}$, $0 < \gamma \leq 1$, $a > 0$. Conditions \mathbf{E}'_γ and $\mathbf{F}'_{a,\gamma}$ are necessary and sufficient for weak convergence $F_\varepsilon(uu_\varepsilon) \Rightarrow F_{a,\gamma}(u)$ as $\varepsilon \rightarrow 0$.*

In conclusion let us give and comment some sufficient conditions providing diffusion and stable approximations for ruin probabilities.

The case $\gamma = 1$ corresponds to so called diffusion approximations of risk processes. The traditional way for obtaining a diffusion type asymptotic is based on approximation of risk process in a proper way by a Wiener process with a shift. Typical conditions assume finiteness of the second moment of the claim distribution $H(t)$ (or equivalently, finiteness of expectation for the steady claim distribution $G(t) = \frac{1}{\mu} \int_0^t (1 - H(s)) ds$):

$$\mathbf{O}:: \mu_2 = \int_0^\infty z^2 H(ds) < \infty.$$

Obviously, condition \mathbf{O} is sufficient for condition \mathbf{E}'_1 to hold. Note however that the necessary and sufficient condition \mathbf{E}'_1 does not require the finiteness of the second moment of the claim distribution.

In this case, condition $\mathbf{F}'_{a,1}$ takes the following simple equivalent form:

$$\mathbf{P}_a:: (1 - \alpha_\varepsilon)u_\varepsilon \rightarrow b = \mu_2/2\mu a \text{ as } \varepsilon \rightarrow 0.$$

As was mentioned in Remark 1, the corresponding limiting distribution is in this case the exponential one, with the parameter a . This is consistent with the classical form of diffusion approximation for ruin probabilities.

Case $\gamma \in (0, 1)$ corresponds to so-called stable approximation for ruin probabilities. Remark 1 explains the sense of using the term “stable”.

Remind that, according Lemma 6, condition \mathbf{E}'_γ is equivalent to the following condition which require regular variation for the tail probabilities of the steady claim distribution $G(t) = \frac{1}{\mu} \int_0^t (1 - H(s)) ds$:

$$\mathbf{R}_\gamma:: \frac{1}{\mu} \int_t^\infty (1 - H(s)) ds \sim \frac{t^{-\gamma} L(t)}{\Gamma(1-\gamma)} \text{ as } t \rightarrow \infty.$$

As follows from theorems about regularly varying functions (see, for example, Feller (1966)), the following condition imposing requirement of regular variation for the tail probabilities of the claim distribution $H(t)$ is sufficient for condition \mathbf{R}_γ to hold:

$$\mathbf{R}'_\gamma:: 1 - H(t) \sim \frac{t^{-(\gamma+1)} L(t) \gamma \mu}{\Gamma(1-\gamma)} \text{ as } t \rightarrow \infty.$$

As far as condition $\mathbf{F}'_{a,\gamma}$ is concerned, it, in this case, can be formulated in the following form equivalent to condition $\mathbf{H}_{a,\gamma}$:

$$\mathbf{S}_{a,\gamma}:: \frac{L(u_\varepsilon)}{(1-\alpha_\varepsilon)u_\varepsilon^\gamma} \rightarrow a \text{ as } \varepsilon \rightarrow 0.$$

REFERENCES

1. Anisimov, V.V., Zakusilo, O.K. and Donchenko, V.S. (1987) Elements of Queuing and Asymptotical Analysis of Systems. Lybid', Kiev
2. Asmussen, S. (1987, 2003) Applied Probability and Queues. Wiley Series in Probability and Mathematical Statistics, Wiley, New York and Applications of Mathematics, **51**, Springer, New York
3. Asmussen, S. (1989) Risk theory in a Markovian environment. Scand. Actuar. J., No. 2, 69–100
4. Asmussen, S. (1996) Rare events in the presence of heavy tails. In: Stochastic Networks, New York, 1995. Lecture Notes in Statistics, **117**, Springer, New York, 197–214

5. Asmussen, S. (2000) Ruin Probabilities. Advanced Series on Statistical Science & Applied Probability, **2**, World Scientific, Singapore
6. von Bahr, B. (1975) Asymptotic ruin probabilities when exponential moments do not exist. *Scand. Actuar. J.*, 6–10
7. Bening, V.E., Korolev, V.Yu. (2002) Generalized Poisson Models and their Applications in Insurance and Finance. Modern Probability and Statistics, VSP, Utrecht
8. Belyaev, Yu.K. (1963) Limit theorems for dissipative flows. *Teor. Veroyatn. Primen.*, **8**, 175–184 (English translation in *Theory Probab. Appl.*, **8**, 165–173)
9. Böker, F., Serfozo, R. (1983) Ordered thinnings of point processes and random measures. *Stoch. Proces. Appl.*, **15**, 113–132
10. Embrechts, P., Goldie, C.M., Veraverbeke, N. (1979) Subexponentiality and infinite divisibility. *Z.Wahrsch. Verm. Geb.*, **49**, 335–347
11. Embrechts, P., Klüppelberg, C., Mikosch, T. (1997) Modelling Extremal Events for Insurance and Finance. Applications of Mathematics, **33**, Springer, Berlin
12. Embrechts, P., Veraverbeke, N. (1982) Estimates for the probability of ruin with special emphasis on the possibility of large claims. *Ins.: Math. and Econ.*, **1**, 55–72
13. Englund, E. (1999a) Perturbed renewal equations with application to M/M queueing systems. 1. *Teor. Ĭmovirn. Mat. Stat.*, **60**, 31–37 (also in *Theory Probab. Math. Statist.*, **60**, 35–42)
14. Englund, E. (1999b) Perturbed renewal equations with application to M/M queueing systems. 2. *Teor. Ĭmovirn. Mat. Stat.*, **61**, 21–32 (also in *Theory Probab. Math. Statist.*, **61**, 21–32)
15. Gasanenko, V.A. (1980) On thinning a semi-Markov process with a countable set of states. *Teor. Veroyatn. Mat. Stat.*, **22**, 25–29 (English translation in *Theory Probab. Math. Statist.*, **22**, 25–29)
16. Gerber, H.U. (1979) An Introduction to Mathematical Risk Theory. Huebner Foundation Monographs, Philadelphia
17. Glynn, P.W. (1990) Diffusion approximations. In: Heyman, D.P., Sobel, M.J. (eds) *Stochastic Models. Handbooks in Operations Research and Management Science*, Vol. 2, North-Holland, Amsterdam, 145–198
18. Gnedenko, B.V. (1964a) On non-loaded duplication. *Izv. Akad. Nauk SSSR, Ser. Tekh. Kibernet.*, No. 4, 3–12
19. Gnedenko, B.V. (1964b) Doubling with renewal. *Izv. Akad. Nauk SSSR, Ser. Tekh. Kibernet.*, No. 5, 111–118
20. Gnedenko, B.V., Fraier, B. (1969) A few remarks on a result by I.N. Kovalenko. *Litov. Mat. Sbornik*, **9**, 463–470
21. Gnedenko, B.V., Korolev, V.Yu. (1996) Random Summation. Limit Theorems and Applications. CRC Press, Boca Raton, FL
22. Gnedenko, B.V., Kovalenko, I.N. (1964, 1987) Introduction to Queueing Theory. Nauka, Moscow (English editions: Israel Program for Scientific Translations, Jerusalem; Daniel Davey & Co., Inc., Hartford, Conn. (1968) and *Mathematical Modeling*, **5**, Birkhäuser, Boston (1989))
23. Grandell, J. (1977) A class of approximations of ruin probabilities. *Scand. Actuar. J., Suppl.*, 37–52
24. Grandell, J. (1991) Aspects of Risk Theory. Probability and its Applications, A Series of the Applied Probability Trust, Springer, New York
25. Gyllenberg, M., Silvestrov, D.S. (1994) Quasi-stationary distributions of a stochastic metapopulation model. *J. Math. Biol.*, **33**, 35–70

26. Gyllenberg, M., Silvestrov, D.S. (2000b) Cramér–Lundberg approximation for nonlinearly perturbed risk processes. *Insurance: Math. Econom.*, **26**, 75–90
27. Harrison, J.M. (1977) Ruin problems with compound assets. *Stoch. Proces. Appl.*, **5**, 67–79
28. Iglehart, D.L. (1969). Diffusion approximation in collective risk theory. *J. Appl. Probab.*, **6**, 285–292
29. Ivchenko, G.I., Kashtanov, V.A., Kovalenko, I.N. (1979) *Theory of Queueing*. Vysshaya Shkola, Moscow
30. Jagers, P. (1974) Aspects of random measures and point processes. In: *Advances in Probability and Related Topics*, Vol. 3. Dekker, New York, 179–239
31. Jagers, P., Lindvall, T. (1974) Thinning and rare events in point processes. *Z. Wahrsch. Verw. Gebiete*, **28**, 89–98
32. Kalashnikov, V.V. (1997) *Geometric Sums: Bounds for Rare Events with Applications*. Mathematics and its Applications, **413**, Kluwer, Dordrecht
33. Kalashnikov, V.V. and Rachev, S.T. (1990) *Mathematical Methods for Construction of Queueing Models*. Wadsworth & Brooks / Cole, Pacific Grove, California.
34. Kalashnikov, V.V., Vsekhsvyatski, S.Y. (1989) On the connection of Rényi's theorem and renewal theory. In: Zolotarev, V.M., Kalashnikov, V.V. (eds) *Stability Problems for Stochastic Models*. Proceedings of the Eleventh International Seminar held in Sukhumi, 1987. *Lecture Notes in Mathematics*, **1412**, Springer, Berlin, 83–102
35. Kallenberg, O. (1975) Limits of compound and thinned point processes. *J. Appl. Probab.*, **12**, 269–278
36. Kaplan, E.I. (1979) Limit theorems for exit times of random sequences with mixing. *Teor. Veroyatn. Mat. Stat.*, **21**, 53–59 (English translation in *Theory Probab. Math. Statist.*, **21**, 59–65)
37. Kartashov, N.V. (1991) Inequalities in Renyi's theorem. *Theory Probab. Math. Statist.*, **45**, 23–28
38. Kingman, J.F. (1963) The exponential decay of Markovian transition probabilities. *Proc. London Math. Soc.*, **13**, 337–358
39. Klüppelberg, C., Stadtmüller, U. (1998) Ruin probability in the presence of heavy-tails and interest rates. *Scand. Act. J.*, **1**, 49–58
40. Korolev, V.Yu. (1989) The asymptotic distributions of random sums. In: Zolotarev, V.M., Kalashnikov, V.V. (eds) *Stability Problems for Stochastic Models*. Proceedings of the Eleventh International Seminar held in Sukhumi, 1987. *Lecture Notes in Mathematics*, **1412**, Springer, Berlin, 110–123
41. Kovalenko, I.N. (1965) On the class of limit distributions for thinning flows of homogeneous events. *Litov. Mat. Sbornik*, **5**, 569–573
42. Kovalenko, I.N. (1975) *Studies in the Analysis of Reliability of Complex Systems*. Naukova Dumka, Kiev
43. Kovalenko, I.N. (1977) Limit theorems of reliability theory. *Kibernetika*, **6**, 106–116 (English translation in *Cybernetics* **13** 902–914 (1977))
44. Kovalenko, I.N. (1980) *Analysis of Rare Events in Estimation of Efficiency and Reliability of Systems*. Sovetskoe Radio, Moscow
45. Kovalenko, I.N. (1994) Rare events in queueing theory – a survey. *Queueing Systems Theory Appl.*, **16**, No. 1-2, 1–49
46. Kovalenko, I.N., Kuznetsov, N.Yu. (1988) *Methods for Calculating Highly Reliable Systems*. Radio and Svyaz', Moscow
47. Kovalenko, I.N. Kuznetsov, N.Yu., Pegg, P.A. (1997) *Mathematical Theory of Reliability of Time Dependent Systems with Practical Applications*, Wiley Series in Probability and Statistics, Wiley, New York

48. Kruglov, V.M., Korolev, V.Yu. (1990) Limit Theorems for Random Sums. Izdatel'stvo Moskovskogo Universiteta, Moscow
49. Lindvall, T. (1976) An invariance principle for thinned random measures. *Studia Sci. Math. Hung.*, **11**, No. 3-4, 269–275
50. Masol, V.I. (1973) On the asymptotic behaviour of the time of arrival at a high level by a regenerating stable process with independent increments. *Teor. Verojatn. Mat. Stat.*, **9**, 131–136 (English translation in *Theory Probab. Math. Statist.*, **9**, 139–144)
51. Melamed, J.A. (1989) Limit theorems in the set-up of summation of a random number of independent identically distributed random variables. In: Zolotarev, V.M., Kalashnikov, V.V. (eds) *Stability Problems for Stochastic Models. Proceedings of the Eleventh International Seminar held in Sukhumi, 1987. Lecture Notes in Mathematics*, **1412**, Springer, Berlin, 194–228
52. Mogyoródi, J. (1971) Some remarks on the rarefaction of the renewal processes. *Litov. Mat. Sbornik*, **11**, 303–315
53. Mogyoródi, J. (1972a) On the rarefaction of renewal processes, I, II. *Studia Sci. Math. Hung.*, **7**, 258–291, 293–305
54. Mogyoródi, J. (1972b) On the rarefaction of renewal processes, III, IV, V, VI. *Studia Sci. Math. Hung.*, **8**, 21–28, 29–38, 193–198, 199–209
55. Råde, L. (1972a) Limit theorems for thinning of renewal point processes. *J. Appl. Probab.*, **9**, 847–851
56. Råde, L. (1972b) *Thinning of Renewal Point Processes. A Flow Graph Study. Matematisk Statistik AB, Göteborg*
57. Rényi, A. (1956) A characterization of Poisson processes. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, **1**, 519–527 (in Hungarian)
58. Schmidli, H. (1992) A General Insurance Risk Model. Ph.D. Thesis, ETH, Zurich
59. Schmidli, H. (1997) An extension to the renewal theorem and an application to risk theory. *Ann. Appl. Probab.*, **7**, 121–133
60. Serfozo, R.F. (1976) Compositions, inverses and thinnings of random measures. *Z. Wahrsch. Verw. Gebiete*, **37**, 253–265
61. Serfozo, R.F. (1984a) Rarefactions of compound point processes. *J. Appl. Probab.*, **21**, 710–719
62. Serfozo, R.F. (1984b) Thinning of cluster processes: Convergence of sums of thinned point processes. *Math. Oper. Res.*, **9**, 522–533
63. Siegmund, D. (1975) The time until ruin in collective risk theory. *Mitt. Verein. Schweiz. Versicherungsmath.*, **75**, No. 2, 157–165
64. Silvestrov, D.S. (1969) Limit theorems for a non-recurrent walk connected with a Markov chain. *Ukr. Mat. Zh.*, **21**, 790–804
65. Silvestrov, D.S. (2000) Perturbed renewal equation and diffusion type approximation for risk processes. *Teor. Īmovirn. Mat. Stat.*, **62**, 134–144 (English translation in *Theory Probab. Math. Statist.*, **62**, 145–156)
66. Silvestrov, D.S., Drozdenko, M.O. (2006) Necessary and sufficient conditions for weak convergence of first-rare-event times for semi-Markov processes. I. *Theory Stoch. Process.* **12(28)**, no.1-2, 161–199
67. Solov'ev, A.D. (1971) Asymptotical behaviour of the first occurrence time of a rare event in a regenerative process. *Izv. Akad. Nauk SSSR, Ser. Tekh. Kibernet.*, No. 6, 79–89 (English translation in *Engrg. Cybernetics*, **9**, No. 6, 1038–1048 (1971))
68. Solov'ev, A.D. (1983) Analytical methods for computing and estimating reliability. In: Gnedenko, B.V. (ed) *Problems of Mathematical Theory of Reliability. Radio i Svyaz'*, Moscow, 9–112

69. Szantai, T. (1971a) On limiting distributions for sums of random variables concerning the rarefaction of recurrent processes. *Studia Sci. Math, Hung.*, **6**, 443–452
70. Szantai, T. (1971b) On an invariance problem related to different rarefactions of recurrent processes. *Studia Sci. Math, Hung.*, **6**, 453–456
71. Szász, D. (1971a) Limit theorems for stochastic processes stopped at random. *Teor. Veroyatn. Primen.*, **16**, 567–569 (English translation in *Theory Probab. Appl.*, **16**, 554–555)
72. Szynal, D. (1976) On limit distribution theorems for sums of a random number of random variables appearing in the study of rarefaction of a recurrent process. *Zastosow. Mat.*, **15**, 277–288
73. Thorin, O., Wikstad, N. (1976) Calculation of ruin probabilities when the claim distributions is lognormal. *Astin Bulletin*, IX, 231–246
74. Tomko, J. (1974) On rarefaction of multivariate point processes. In: *Progress in Statistics*, Vol. II. Coll. Math. Soc. Janos Bolyai, **9**, 843–868
75. Zinchenko, N. (1999) Risk processes in the presence of large claims. In: Silvestrov, D., Yadrenko, M., Borysenko, O., Zinchenko, N. (eds) *Proceedings of the Second International School on Applied Statistics, Financial and Actuarial Mathematics*, Kyiv, 1999. *Theory Stoch. Proces.*, **5(21)**, No. 1-2, 200–211

DEPARTMENT OF MATHEMATICS AND PHYSICS, MÄLARDALEN UNIVERSITY, BOX 883, SE-721 23 VÄSTERÅS, SWEDEN

E-mail address: dmitrii.silvestrov@mdh.se

E-mail address: myroslav.drozdenko@mdh.se